# **Popular Transdermal Preparations**

# **Rheumatoid Arthritis/Joint Pain**

Ketorolac 8%, DMSO 10 -20%, Tetracaine 2-5%, + CMO 2-6%, in Lipoderm

Diclofenac 8-12%, DMSO 10-20%, Tetracaine 2-5%, + CMO 2-10% in Lipoderm

Note: Use ketoprofen 20% if a rash develops from diclofenac

Note: Ketorolac PO to be taken 30 days maximum. If used on the knee, there is less than 5% ending up in the systemic circulation so it can be taken longer. Not to be used for non-joint areas.

### **Lower Back Pain**

Diclofenac 8-12%, Baclofen 2-5%, DMSO 5-10%, Tetracaine 2-5%, + CMO 2-6% in Lipoderm

## **Post-Herpetic Neuralgia**

Ketamine 10-15%, Morphine 0.5-2%, Amitriptyline 2-5%, Lidocaine 2-5% in Lipoderm

Use the following spray if the skin is too sensitive to apply the cream Benzocaine 5%, Lidocaine 10%, Tetracaine 5% Spray

# **Neuropathic Pain**

Loperamide\* 5-10%, Gabapentin 6-10%, Amitriptyline 2-5%, Lidocaine 2-5% in Lipoderm \*Note: Loperamide is a weak mu agonist and extremely expensive.

Ketamine 5-15%, Morphine 0.5-1%, Gabapentin 6-10%, Lidocaine 2-5% in Lipoderm +Amitriptyline 2-5%,

#### **Radiation Burns**

Ketamine 2.5%, Morphine 0.5%, Lidocaine 2% + Gabapentin 6% in Lipoderm

Note: Skin has increased absorptive abilities so use lower concentrations to reduce systemic drug levels

## Raynaud's Syndrome

Nifedipine 2-15%, Sildenafil 2-5%, Pentoxyphylline 5-15% in Lipoderm

Nitroglycerin\* 0.9% in Glaxal Base

Note: Nitroglycerin will probably cause a headache when 1<sup>st</sup> starting treatment but will disappear. Start with once a day applications to build tolerance.

### **Radiation Proctitis**

Sucralfate 2-3g/15-20mL enema PR OD - BID x 12 days

Rectal Bleeding? Try 4g/20mL

May need to lower volume if urge to defecate present when using enema Think of Glutamine supplementation orally – it is the building block of the GI Tract!

## Transmucosal Pain – Rectal and Vaginal

Morphine 5-20mg, lidocaine 10-50mg, sucralfate 40-80mg in **MBK** suppositories Can add ketamine 5-20mg

### Suppositories seem to last longer than cream

Morphine 0.5-2%, lidocaine 2-5%, sucralfate 4-8% in Emollient Cream Can add ketamine 5-15%

Clotrimazole 2%, lidocaine 2-5%, sucralfate 4-8% in Versabase

Ketamine 2.5-5%, hydromorphone 0.5-2%, lidocaine 2-5%, sucralfate 4-8% in Versabase

Ketamine 2.5-5%, morphine 0.5-2%, lidocaine 2-5%, sucralfate 4-8% in Versabase

Lidocaine 2-5%, sucralfate 4-8% in Versabase

\*Note: Versabase has better penetration than Emollient Cream

# **Chemo Induced Peripheral Neuropathy**

Ketamine 5-15%, baclofen 2-5%, lidocaine 2-5% <u>+</u> morphine 0.5-2% in Lipoderm